

Human Recombinant Gonadotrophin Releasing Hormone Receptor Stable Cell Line

Cat. No. M00426

Version 05282014

I	Introduction	1
II	Background.....	1
III	Representative Data.....	2
IV	Thawing and Subculturing.....	2
V	References.....	3
	Limited Use License Agreement.....	4

I. INTRODUCTION

Catalog Number: M00426

Cell Line Name: CHO-K1/GNRHR/G α 15

Gene Synonyms: GNRHR; GNRHR1; GRHR; LHRHR; LRHR

Expressed Gene: Genbank Accession Number NM_000406; no expressed tags

Host Cell: CHO-K1/G α 15Quantity: Two vials of frozen cells (3×10^6 per vial)

Stability: 16 passages

Application: Functional assay for GNRHR receptor

Freeze Medium: 45% culture medium, 45% FBS, 10% DMSO

Complete Growth Medium: Ham's F12, 10% FBS

Culture Medium: Ham's F12, 10% FBS, 400 μ g/ml G418, 100 μ g/ml HygromycinB

Mycoplasma Status: Negative

Storage: Liquid nitrogen immediately upon delivery

II. BACKGROUND

The gonadotropin-releasing hormone receptor (GNRHR), also known as the luteinizing hormone releasing hormone receptor (LHRHR), is a member of the seven-transmembrane, G-protein coupled receptor (GPCR) family. It is expressed on the surface of pituitary gonadotrope cells as well as lymphocytes, breast, ovary, and prostate. Upon activation, the LHRHR stimulates tyrosine phosphatase and elicits the release of LH from the pituitary. Evidence exists showing the presence of LHRH and its receptor in extrapituitary tissues as well as a role in progression of some cancers.

III. REPRESENTATIVE DATA

Concentration-dependent stimulation of intracellular calcium mobilization by LH-RH in CHO-K1/GNRHR/Gα15 and CHO-K1/Gα15 cells




Figure 1. LH-RH-induced concentration-dependent stimulation of intracellular calcium mobilization in CHO-K1/GNRHR/Gα15 and CHO-K1/Gα15 cells. The cells were loaded with Calcium-4 prior to stimulation with a GNRHR receptor agonist, LH-RH. The intracellular calcium change was measured by FlexStation. The relative fluorescent units (RFU) were plotted against the log of the cumulative doses (10-fold dilution) of LH-RH (Mean ± SD, n = 2). The EC₅₀ of LH-RH on GNRHR co-expressing with Gα15 in CHO-K1 cells was 41 nM. The S/B of LH-RH on GNRHR co-expressing with Gα15 in CHO-K1 cells was 16.

Notes:

- EC₅₀ value is calculated with four parameter logistic equation:

$$Y = \text{Bottom} + \frac{(\text{Top} - \text{Bottom})}{(1 + 10^{-(\text{LogEC}_{50} - X) \cdot \text{HillSlope}})}$$

X is the logarithm of concentration. Y is the response
 Y is RFU and starts at Bottom and goes to Top with a sigmoid shape.
- Signal to background Ratio (S/B) = Top/Bottom

IV. THAWING AND SUBCULTURING

Thawing Protocol

- Remove the vial from liquid nitrogen tank and thaw cells quickly in a 37°C water-bath.
- Just before the cells are completely thawed, decontaminate the outside of the vial with 70% ethanol and transfer the cells to a 15 ml centrifuge tube containing 9 ml of complete growth medium.
- Pellet cells by centrifugation at 200 x g force for 5 min, and remove the medium.
- Resuspend the cells in complete growth medium.
- Transfer the cell suspension to a 10 cm dish with 10 ml of complete growth medium.
- Grow the cells in incubator with 37°C, 5 %CO₂.
- In the following day, replace the cells with fresh medium contains antibiotic.

Sub-culturing Protocol

1. Remove the culture medium from cells.
2. Wash cells with PBS (pH=7.4) to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 ml of 0.05% (w/v) Trypsin- EDTA (GIBCO, Cat No. 25300) solution into 10 cm dish and observe the cells under an inverted microscope until cell layer is dispersed (usually within 3 to 5 minutes).
Note: To avoid cells clumping, do not agitate the cells by hitting or shaking the dish while waiting for the cells detach. If cells are difficult to detach, please place the dish in 37°C incubator for ~2 min.
4. Add 6.0 to 8.0 ml of complete growth medium into dish and aspirate cells by gently pipetting.
5. Centrifuge the cells at 200 x g force for 5min, and remove the medium.
6. Resuspend the cells in culture medium and add the cells suspension to new culture dish.
7. Grow the cells in incubator with 37°C, 5 %CO₂.

Subcultivation Ratio: 1:3 to 1:8 weekly.

Medium Renewal: Every 2 to 3 days

V. REFERENCES

1. Millar RP, (2005) GnRHs and GnRH receptors. *Anim. Reprod. Sci.* 88(1-2): 5–28.
2. Harrison GS, Wierman ME, Nett TM, *et al.* (2004) Gonadotropin-releasing hormone and its receptor in normal and malignant cells. *Endocr. Relat. Cancer.* 11(4): 725–48.

GenScript USA Inc,

860 Centennial Ave.

Piscataway, NJ 08854

Toll-Free: 1-877-436-7274

Tel: 1-732-885-9188, Fax: 1-732-210-0262

Email: product@genscript.com

Web: <http://www.genscript.com>

For Research Use Only.

860 Centennial Ave., Piscataway, NJ 08854, USA

Toll-Free: 1-877-436-7274 Tel: 1-732-885-9188 Fax: 1-732-210-0262 Email: product@genscript.com Web: www.genscript.com

Limited Use License Agreement

This is a legal agreement between you (Licensee) and GenScript USA Inc. governing use of GenScript's stable cell line products and protocols provided to licensee. By purchasing and using the stable cell line, the buyer agrees to comply with the following terms and conditions of this label license and recognizes and agrees to such restrictions:

- 1) The products are not transferable and will be used at the site where they were purchased. Transfer to another site owned by buyer will be permitted only upon written request by buyer followed by subsequent written approval by GenScript.
- 2) The purchaser cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party.
- 3) The products sold by GenScript are for laboratory and animal research purposes only. The products are not to be used on humans, for consumption, or for any unlawful uses.

GenScript USA Inc. will not assert against the buyer a claim of infringement of patents owned or controlled by GenScript USA Inc. and claiming this product based upon the manufacture, use or sale of a clinical diagnostic, therapeutic and vaccine, or prophylactic product developed in research by the buyer in which this product or its components has been employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on the use of this product for other purposes, contact Marketing Department, GenScript USA Inc., 120 Centennial Avenue, Piscataway, New Jersey 08840, U.S.A. Phone: 1-732-885-9188. Fax: 1-732-210-0262. Email: marketing@genscript.com.